

NCI-H295R Cells | 300483

Renseignements généraux

Description H295R was adapted from the NCI-H295 pluripotent adrenocortical carcinoma cell line established by A.F. Gazdar and associates (1990) from a carcinoma of the adrenal cortex. The original cells were adapted to a culture medium which decreased the population doubling time from 5 days to 2 days. The adapted cells were selected to grow in a monolayer, in contrast to the original cells which grew in suspension. This cell line retains the ability to produce adrenal androgens. It is responsive to angiotensin II and potassium ions.

Organism Human

Tissue Adrenal gland

Disease Carcinoma

Synonyms NCI-H295R, NCI H295R, NCIH295R, H-295R, H295R-S1

Caractéristiques

Age 48 years

Gender Female

Ethnicity Caucasian

Morphology Epithelial-like

Growth properties Monolayer, adherent

Données réglementaires

Citation NCI-H295R (Cytion catalog number 300483)

Biosafety level 1

NCBI_TaxID 9606

CellosaurusAccession CVCL_0458

Données biomoléculaires

NCI-H295R Cells | 300483

Products Aldosterone, cortisol, C19 steroids

Manipulation

Culture Medium You can purchase our ready-to-use NCI-H295R Cell Growth Medium (820402) or choose to supplement DMEM:Ham's F12 (1:1), w: 3.1 g/L Glucose, w: 2.5 mM L-Glutamine, w: 15 mM HEPES, w: 0.5 mM Sodium pyruvate, w: 1.2 g/L NaHCO₃ (Cytion article number 820400a) with the below additives

Supplements Supplement the medium with 5% FBS, 0.00625 mg/mL insulin, 0.00625 mg/mL transferrin, 6.25 ng/mL selenium, 1.25 mg/mL bovine serum albumin, 0.00535 mg/mL linoleic acid

Dissociation Reagent Accutase

Subculturing Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

Fluid renewal 2 to 3 times per week

Post-Thaw Recovery 48 hours

Freeze medium As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

NCI-H295R Cells | 300483

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $300 \times g$ for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C , 5% CO_2 , humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

Contrôle de la qualité et analyse moléculaire

NCI-H295R Cells | 300483

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.