

XXXXXXXXXXXXXXXXXXXX

XIX XIX XIX XIX XIX XIX XIX XIX

~~XXXXXXXXXX XIXX-XXXXXXXXXXXXXXXXXX~~

XXXXXX

HEK293T BHK-21 | 305252

Supplements 10% FBS 1% NEAA

Dissociation Reagent Trypsin

Subculturing HEK293T cells are grown in DMEM supplemented with 10% FBS and 1% NEAA. Cells are passaged when they reach 80-90% confluency.

Seeding density 1 to 3 x 10⁴ cells/cm²

Fluid renewal 2 to 3 times per week

Freeze medium HEK293T cells are frozen in DMEM supplemented with 10% FBS and 10% DMSO. Cells are frozen in 1 mL aliquots and stored at -80°C.

Thawing and Culturing Cells

1. HEK293T cells are thawed in a water bath at 37°C. Cells are then centrifuged at 300 x g for 5 minutes and the supernatant is removed.
2. HEK293T cells are resuspended in DMEM supplemented with 10% FBS and 1% NEAA. Cells are then seeded into a 25 cm² flask.
3. HEK293T cells are grown in DMEM supplemented with 10% FBS and 1% NEAA. Cells are passaged when they reach 80-90% confluency.
4. HEK293T cells are grown in DMEM supplemented with 10% FBS and 1% NEAA. Cells are passaged when they reach 80-90% confluency.
5. HEK293T cells are grown in DMEM supplemented with 10% FBS and 1% NEAA. Cells are passaged when they reach 80-90% confluency.
6. HEK293T cells are grown in DMEM supplemented with 10% FBS and 1% NEAA. Cells are passaged when they reach 80-90% confluency.
7. HEK293T cells are grown in DMEM supplemented with 10% FBS and 1% NEAA. Cells are passaged when they reach 80-90% confluency.
8. HEK293T cells are grown in DMEM supplemented with 10% FBS and 1% NEAA. Cells are passaged when they reach 80-90% confluency.

Incubation Atmosphere 37°C, 5% CO₂, humidified

Shipping Conditions HEK293T cells are shipped in DMEM supplemented with 10% FBS and 1% NEAA. Cells are shipped at 4°C.

Storage Conditions HEK293T cells are stored in DMEM supplemented with 10% FBS and 1% NEAA. Cells are stored at -80°C.

Product sheet

BHK-21 | 305252

PCR

Sterility

The BHK-21 is a PCR tube that is designed for use in PCR applications. The tube is made of polypropylene and is designed to be used in a PCR thermocycler. The tube is designed to be used in a PCR thermocycler and is designed to be used in a PCR thermocycler.