

HEK293-TACD2 Cells | 305424

General information

Description

Disclaimer: The prices displayed for cell lines are exclusively for academic/not-for-profit customers. For commercial entities the price is approximately €6,250.

If you represent a commercial entity or are unsure which category applies, please [contact us](#).

The HEK293-TACD2 cell line is a stable recombinant HEK293 cell line engineered to express the TACD2 receptor at a medium-high level, approximately 10,000 molecules per cell. This cell line was developed using inscreenex's landing pad technology, which ensures precise and reproducible integration of the TACD2 gene at a specific, pre-validated genomic locus. TACD2, also known as TROP2 or GA733-1, is a tumor-associated calcium signal transducer that plays a key role in intracellular calcium signaling, crucial for cellular processes such as growth, division, and differentiation. Overexpression of TACD2 has been observed in various carcinomas, including colorectal, gastric, and pancreatic cancers, making it a significant target for antibody-drug conjugates and immunotherapy.

The expression of TACD2 in this cell line was confirmed using flow cytometry with a target-specific antibody, ensuring reliable and consistent receptor density across the cell population.

Organism

Human

Tissue

Fetal Kidney

Disease

Transformed/immortalized; non-tumorigenic (HEK293 background)

Applications

TROP2 (TACD2/TACSTD2)-targeted antibody and ADC development (sacituzumab govitecan analogs); triple-negative breast, lung and urothelial cancer research; ADCC/CDC assays; flow cytometry

Characteristics

Age

Fetus

Gender

Female

Morphology

Epithelial-like

Cell type

Epithelial cells

Growth properties

Monolayer, adherent

Regulatory Data

HEK293-TACD2 Cells | 305424**Citation** HEK293-TACD2 (Cytion catalog number 305424)**Biosafety level** 1**NCBI_TaxID** 9606**CellosaurusAccession** CVCL_6G29**GMO Status** GMO-S1: This HEK293 line contains a TACD2 expression construct for receptor-binding and functional analyses. This classification applies only within Germany and may differ elsewhere.**Biomolecular Data****Receptors expressed** TACD2 (TROP2 or GA733-1)**Handling****Culture Medium** RPMI 1640, w: 2.0 mM stable Glutamine, w: 2.0 g/L NaHCO₃ (Cytion article number 820700a)**Supplements** Supplement the medium with 10% FBS, 1 mM sodium pyruvate, 10 mM HEPES, 1% NEAA. Add Geneticin (G418-Sulfat) to achieve a final concentration of 1 mg/mL.**Dissociation Reagent** Trypsin-EDTA**Doubling time** approx. 24-36 hours**Subculturing** For routine adherent cell culture: Aspirate the old culture medium from the adherent cells, and wash them with PBS to remove any remaining medium. After aspirating the PBS, add the appropriate volume of Trypsin/EDTA solution based on the culture vessel size (e.g., 1 ml for a T25 flask, 3 ml for a T75 flask) and incubate at room temperature or 37°C until the cells detach (5-10 minutes). Monitor detachment under a microscope, and gently tap the vessel if necessary to release the cells. Once detached, add complete medium to inactivate the Trypsin/EDTA, gently resuspend the cells, and transfer an aliquot of the cell suspension into a new culture vessel containing fresh medium. Place the vessel in an incubator set to 37°C with 5% CO₂, and change the medium every 2-3 days.**Split ratio** 1 to 5**Seeding density** 2 to 4 x 10⁴ cells/cm²

HEK293-TACD2 Cells | 305424

Fluid renewal 2 to 3 times per week

Post-Thaw Recovery

After thawing, split the cells at a ratio of 1:2 to 1:3 in T25 flasks and allow the cells to recover from the freezing process and to adhere for at least 24 hours.

For best attachment and viability after thawing the cells, we recommend using Collagen-coated flasks or plates for the initial seeding after cryo-recovery. Collagen coating is not required for subsequent routine culture of the cells.

Freeze medium

As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C, 5% CO₂, humidified atmosphere.

HEK293-TACD2 Cells | 305424

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.