

SHP-77 Cells | 305498

General information

Description

The SHP-77 cell line is a human small-cell lung carcinoma (SCLC) model. It was derived from a primary lung tumor and is used extensively in cancer research, particularly for studies focused on lung cancer biology and drug development. SHP-77 cells exhibit the classic characteristics of SCLC, including rapid growth and high tumorigenic potential in xenograft models. This cell line is known for its ability to proliferate in serum-supplemented culture media and has been utilized in various experimental setups, such as studies of oncogenic signaling pathways and therapeutic response to chemotherapeutic agents.

SHP-77 cells are part of the Cancer Cell Line Encyclopedia (CCLE), a resource that enables researchers to correlate genetic profiles with drug sensitivities. Genomic profiling of SHP-77 has revealed mutations and alterations in critical oncogenes and tumor suppressors, providing a platform for studying the molecular mechanisms underlying SCLC pathogenesis. The cell line has also been included in drug screening studies, offering insights into its pharmacological vulnerabilities and aiding in the identification of compounds with therapeutic potential for lung cancer.

Organism

Human

Tissue

Lung, left upper lobe

Disease

small cell carcinoma

Applications

3D cell culture, Cancer research

Synonyms

SHP77, Shadyside Hospital Pittsburgh-77

Characteristics

Age

54 years

Gender

Male

Ethnicity

Caucasian

Morphology

Round cells

Cell type

Epithelial cells

Growth properties

Mixed: suspension with some loosely adherent cells

Regulatory Data

SHP-77 Cells | 305498

Citation	SHP-77 (Cytion catalog number 305498)
----------	---------------------------------------

Biosafety level	1
-----------------	---

NCBI_TaxID	9606
------------	------

CellosaurusAccession	CVCL_1693
----------------------	-----------

Biomolecular Data

Antigen expression	Blood Type O; Rh +; CD56; CD57 (HNK-1,Leu-7)
--------------------	--

Tumorigenic	Yes; Yes, the cells form tumors in athymic nude mice, and usually grow as circumscribed nodules without evidence of metastases
-------------	--

Mutational profile	Mutation: ABL1, Simple, p.Val1128Glu (c.3383T>A), Zygosity=Heterozygous; Mutation: KRAS, Simple, p.Gly12Val (c.35G>T), Homozygous; Mutation: RAC1, Simple, p.Tyr32Cys (c.95A>G), Heterozygous; Mutation: TP53, Simple, p.Cys176Trp (c.528C>G), Homozygous
--------------------	---

Handling

Culture Medium	RPMI 1640, w: 2.0 mM stable Glutamine, w: 2.0 g/L NaHCO ₃ (Cytion article number 820700a)
----------------	--

Supplements	Supplement the medium with 10% FBS
-------------	------------------------------------

Doubling time	85 hours
---------------	----------

Fluid renewal	2 to 3 times per week
---------------	-----------------------

Freeze medium	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.
---------------	---

SHP-77 Cells | 305498

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C, 5% CO₂, humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196 °C. Storage at -80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis

SHP-77 Cells | 305498

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.