

A2780-GFP | 305676

General information

Description

A2780-GFP is a genetically modified derivative of the human ovarian carcinoma cell line A2780, engineered to stably express green fluorescent protein (GFP). The parental A2780 cell line was established from an ovarian tumor in an adult patient and is widely used as a model for epithelial ovarian cancer, particularly in studies of chemotherapy response. It is known for its relative sensitivity to platinum-based agents such as cisplatin, making it a valuable system for investigating mechanisms of drug action and resistance. The GFP-expressing variant preserves these biological characteristics while incorporating a fluorescent reporter for enhanced experimental utility.

The stable expression of GFP enables real-time visualization and quantitative analysis of A2780-GFP cells in a variety of experimental settings. Fluorescence-based detection allows researchers to monitor cell proliferation, morphology, migration, and viability in vitro, as well as to track tumor growth and dissemination in vivo. This is particularly advantageous in xenograft and metastasis models, where GFP facilitates discrimination of tumor cells from surrounding host tissue. The fluorescent signal is generally stable across passages, although expression levels may vary depending on the transduction method and clonal selection.

A2780-GFP retains the core molecular and phenotypic features of the parental A2780 line, including pathways associated with DNA damage response, apoptosis, and chemotherapeutic sensitivity. As such, it is commonly used in high-content imaging assays, drug screening platforms, and co-culture systems where spatial and temporal resolution of tumor cell behavior is required. The addition of GFP significantly enhances the versatility of this model, supporting applications in ovarian cancer research, therapeutic evaluation, and studies of tumor cell dynamics.

Organism Human

Tissue Ovary

Disease Ovarian endometrioid adenocarcinoma

Characteristics

Age Age unspecified

Gender Female

Ethnicity African American

Growth properties Adherent

Regulatory Data

A2780-GFP | 305676

Citation A2780-GFP (Cytion catalog number 305676)

Biosafety level 1

NCBI_TaxID 9606

GMO Status GMO-S1: This human A2780 ovarian cancer line contains a lentiviral GFP expression construct for fluorescence-based imaging. This classification applies only within Germany and may differ elsewhere.

Biomolecular Data

Antigen expression GFP

Handling

Culture Medium RPMI 1640, w: 2.0 mM stable Glutamine, w: 2.0 g/L NaHCO₃ (Cytion article number 820700a)

Supplements Supplement the medium with 10% FBS

Dissociation Reagent Accutase

Freeze medium As a cryopreservation medium, we use complete growth medium + 10% DMSO for adequate post-thaw viability.

A2780-GFP | 305676

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $200 \times g$ for 5 minutes, carefully discard the supernatant containing freezing medium.
7. Follow the procedure described under Post-Thaw Recovery

Incubation Atmosphere

37°C , 5% CO_2 , humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis