

## SCLC-22H Cells | 300445

### General information

#### Description

The SCLC-22H cell line was established from the pericardial effusion of a male patient diagnosed with small cell lung cancer (SCLC) of the oat cell type, an aggressive subtype of lung cancer. The SCLC-22H cell line, derived from a small cell lung cancer (SCLC) patient, exhibits a mixture of features typical of both the classic and variant types of SCLC. This intermediate nature makes it a valuable model for studying the transition between these two subtypes. The cell line shows morphological characteristics such as small and large cell-like features, which are typically seen in both small cell and large cell lung cancer, especially when examined in xenografts.

SCLC-22H expresses several neuroendocrine markers, including neuron-specific enolase (NSE), carcinoembryonic antigen (CEA), bombesin, and creatine kinase-BB (CK-BB), which are hallmarks of classic SCLC. However, compared to the closely related SCLC-21H cell line, SCLC-22H has a slower population doubling time and a lower colony-forming efficiency. These biochemical and kinetic properties distinguish it from SCLC-21H, which displays more features of the variant subtype with predominantly large cell morphology.

SCLC-22H is considered an important model for understanding the *in vivo* progression from classic to variant SCLC. Its mixed phenotype suggests that it represents an intermediate or transitional phase, offering insights into how treatment resistance and changes in cell morphology and growth characteristics develop in aggressive lung cancers.

**Organism** Human

**Tissue** Lung

**Disease** Small cell carcinoma

**Metastatic site** Pericardial effusion

**Synonyms** SCLC22H

### Characteristics

**Age** 46 years

**Gender** Male

**Ethnicity** Caucasian

**Morphology** Floating cell aggregates, few single cells

**Growth properties** Suspension

## SCLC-22H Cells | 300445

## Regulatory Data

<b>Citation</b>	SCLC-22H (Cytion catalog number 300445)
<b>Biosafety level</b>	1
<b>NCBI_TaxID</b>	9606
<b>CellosaurusAccession</b>	CVCL_2186

## Biomolecular Data

<b>Tumorigenic</b>	Yes, in nude mice
<b>Reverse transcriptase</b>	Negative
<b>Karyotype</b>	Modal number 43

## Handling

<b>Culture Medium</b>	RPMI 1640, w: 2.0 mM stable Glutamine, w: 2.0 g/L NaHCO <sub>3</sub> (Cytion article number 820700a)
<b>Supplements</b>	Supplement the medium with 10% FBS
<b>Subculturing</b>	Maintain cultures by periodically adding or replacing the medium. Initiate cultures with a density of $5 \times 10^5$ cells/ml and keep the cell concentration within the range of $1 \times 10^5$ to $1 \times 10^6$ cells/ml for optimal growth.
<b>Seeding density</b>	$1 \times 10^5$ cells/ml
<b>Fluid renewal</b>	1 to 2 times per week
<b>Freeze medium</b>	As a cryopreservation medium, we use 50% basal medium + 40% FBS + 10% DMSO, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

## SCLC-22H Cells | 300445

### Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below  $-150^{\circ}\text{C}$  to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a  $37^{\circ}\text{C}$  water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at  $300 \times g$  for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

### Incubation Atmosphere

$37^{\circ}\text{C}$ , 5%  $\text{CO}_2$ , humidified atmosphere.

### Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately  $-78^{\circ}\text{C}$  throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

### Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about  $-150$  to  $-196^{\circ}\text{C}$ . Storage at  $-80^{\circ}\text{C}$  is acceptable only as a short interim step before transfer to liquid nitrogen.

## Quality Control & Molecular Analysis

**SCLC-22H Cells | 300445**

**Sterility**

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.