

HEC-1-A Cells | 305077

General information

Description

HEC-1-A cells are a well-characterized human endometrial adenocarcinoma cell line derived from the malignant tissue of a 71-year-old Caucasian woman. This cell line, established in the mid-1970s, is extensively used in gynecological cancer research, particularly for studying endometrial carcinoma.

Morphologically, HEC-1-A cells are epithelial-like and form a monolayer of polygonal cells when cultured. They exhibit a robust and adherent growth pattern, which is typical of epithelial cells originating from solid tumors. The morphological characteristics of HEC-1-A cells make them a valuable model for studying cellular behaviors that are central to cancer progression, such as adhesion, migration, and invasion.

Genotypically, HEC-1-A cells harbor several genetic aberrations that are relevant to cancer biology, including mutations in key regulatory genes like p53 and PTEN, both of which are commonly mutated in endometrial cancer. These genetic features contribute to the cells' utility in researching the molecular underpinnings of endometrial carcinogenesis and the cellular pathways leading to tumor growth and resistance to therapy.

Research using HEC-1-A cells has significantly advanced our understanding of endometrial cancer, particularly in terms of hormonal influences, genetic mutations, and responses to chemotherapeutic agents. As a result, this cell line continues to be instrumental in developing more effective diagnostic and therapeutic strategies for endometrial carcinoma.

Organism Human

Tissue Uterus, endometrium

Disease Endometrial adenocarcinoma

Synonyms Hec-1-A, HEC-1A, HEC1-A, HEC1A, Hec1A

Characteristics

Age 71 years

Gender Female

Ethnicity Asian

Morphology Epithelial

Growth properties Adherent

Regulatory Data

HEC-1-A Cells | 305077**Citation** HEC-1-A (Cytion catalog number 305077)**Biosafety level** 1**NCBI_TaxID** 9606**CellosaurusAccession** CVCL_0293**Biomolecular Data****Receptors expressed** Receptor expression: platelet activating factor(PAF)**Protein expression** Oncogenes: C-Fos**Antigen expression** Blood Type B, Rh**Tumorigenic** Yes**Handling****Culture Medium** McCoys 5a, w: 3.0 g/L Glucose, w: stable Glutamine, w: 2.0 mM Sodium pyruvate, w: 2.2 g/L NaHCO₃ (Cytion article number 820200a)**Supplements** Supplement the medium with 10% FBS**Dissociation Reagent** Accutase**Subculturing** Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.**Fluid renewal** 2 to 3 times per week**Freeze medium** As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

HEC-1-A Cells | 305077

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $300 \times g$ for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C , 5% CO_2 , humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis

HEC-1-A Cells | 305077

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.