

HROC277 T0 M1 Cells | 300834**General information**

Description	This is one cell line of a series of tumor cell lines which have been established by PD Dr. Michael Linnebacher from Primary CRC resection specimens since 2006.
Organism	Human
Tissue	Colon coecum, UICC IV, Established from a patient-derived xenograft of primary CRC tissue (Colon ascendens, TNM stage T4N0M1R0L0V0, grade G2, Lk(n) +0, Σ Lk(n) 12)
Disease	Adenocarcinoma
Synonyms	HROC277

Characteristics

Age	77 years
Gender	Male
Ethnicity	Caucasian
Morphology	Epithelial-like
Growth properties	Adherent

Regulatory Data

Citation	HROC277 T0 M1 (Cytion catalog number 300834)
Biosafety level	1
NCBI_TaxID	9606
CellosaurusAccession	CVCL_1U86

Biomolecular Data

Protein expression	PTEN-
---------------------------	-------

HROC277 T0 M1 Cells | 300834

Tumorigenic Yes, in immune-suppressed nude mice

Viruses Free of human pathogenic viruses SV40, JC/BK, HBV, HCV, HIV.

MSI-status MSS

Mutational profile K-RasG12A , N-Raswt, H-Raswt, PIK3CAwt, B-Rafwt

Handling

Culture Medium DMEM:Ham's F12 (1:1), w: 3.1 g/L Glucose, w: 2.5 mM L-Glutamine, w: 15 mM HEPES, w: 0.5 mM Sodium pyruvate, w: 1.2 g/L NaHCO₃ (Cytion article number 820400a)

Supplements Supplement the medium with 10% FBS

Dissociation Reagent Accutase

Doubling time 29 hours

Subculturing Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

Seeding density 2×10^4 cells/cm²

Fluid renewal Every 3 to 5 days

Post-Thaw Recovery Few days

Freeze medium As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

HROC277 T0 M1 Cells | 300834

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $300 \times g$ for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C , 5% CO_2 , humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis

HROC277 T0 M1 Cells | 300834

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.