

B-LCL-CDG2 Cells | 302013**General information**

Description	B-LCL-CDG2 is an EBV-transformed B lymphocyte cell line derived from a young girl suffering from PMM2-CDG. PMM2-CDG is a rare inborn error of metabolism, which results in defective synthesis of glycosylated oligosaccharide chains of many tissue and blood glycoproteins and/or glycosphingolipids. The primary cause of defective glycosylation is based on mutations in the enzyme phosphomannomutase 2 (PMM2). There are two distinct mutations for the PMM2 gene.
Organism	Human
Tissue	Peripheral blood
Disease	Congenital Disorders of Glycosylation
Applications	Genotyping of CDG effects in immune cells, functional testing (e.g. B cell surface antigens), testing of cytotoxic drugs, mutational analysis, analysis of apoptotic mechanisms, HLA-typing, impact of defective glycosylation of distinct cellular glycoproteins on diverse functions.

Characteristics

Age	Child
Gender	Female
Ethnicity	Caucasian
Morphology	Round cells
Cell type	B lymphocyte
Growth properties	Suspension, Cluster

Regulatory Data

Citation	B-LCL-CDG2 (Cytion catalog number 302013)
Biosafety level	2
NCBI_TaxID	9606

B-LCL-CDG2 Cells | 302013**CellosaurusAccession** CVCL_A9Y1**Biomolecular Data****Surface antigens** CD60a- (GD3), CD60c-(7-O-acetylated GD3), CD75s+ sialylated lactosaminyl Noligosaccharides), CD77- (Gb3, globotriaosylceramide)**Antigen expression** CD10-, CD19+, CD20+, CD21+, CD22+, CD23+, CD24+, CD37+m CD38+, CD39+, CD40+, CD53+, CD71+, CD72(+), CD73+, CD74 (+), CD80+, CD81+, CD82+, CD83-, CD84-, CD85+, CD86+, MHC class I+, MHC class II+**Viruses** Transformant: EBV**Handling****Culture Medium** RPMI 1640, w: 2.0 mM stable Glutamine, w: 2.0 g/L NaHCO₃ (Cytion article number 820700a)**Supplements** Supplement the medium with 10% heat-inactivated FBS**Subculturing** Maintain cultures by periodically adding or replacing the medium. Initiate cultures with a density of 2×10^5 cells/ml and keep the cell concentration within the range of 1×10^5 to 5×10^5 cells/ml for optimal growth.**Fluid renewal** Once the medium colour turned into yellow**Post-Thaw Recovery** Medium**Freeze medium** As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

B-LCL-CDG2 Cells | 302013

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C, 5% CO₂, humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196 °C. Storage at -80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis

B-LCL-CDG2 Cells | 302013

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.