

## Caco-2 Cells | 300137

### General information

#### Description

Caco-2 cells serve as an advanced in vitro model for the human intestinal barrier, primarily due to their differentiation into a cell monolayer that closely resembles the enterocytes lining the small intestine. When culturing the Caco2 cell line on tissue culture filter inserts with polycarbonate filters, Caco-2 cells undergo spontaneous differentiation. The differentiation of Caco2 cells results in the expression of specialized cell types, complete with microvilli, enzymes, and transporters, paralleling the complex features and mechanisms found in an in vivo situation.

In the context of intestinal absorption studies models, Caco-2 cells, which were derived from a human colorectal adenocarcinoma patient, are instrumental due to their ability to develop high TEER values, signifying intact tight junctions and epithelial barrier function. These properties are crucial for assays like the cholesterol efflux assay and investigations into cellular transport, including the movement of lipid nanoparticles and the detection of protein interactions.

Caco-2 cells are pivotal for intestinal absorption studies, providing a reliable in vitro approximation of the intestinal epithelium. Mimicking intestinal enterocytes, these cells facilitate analyses of oral drug absorption by simulating the intestinal barrier. Researchers utilize Caco-2 cells to predict how substances traverse the intestinal mucosa, which is essential for the pharmacokinetic profiling of oral medications. Furthermore, they are a key tool in investigating intestinal cholesterol uptake, homeostasis and transport, which are vital processes for understanding lipid metabolism and associated diseases.

Caco-2 cells remain a cornerstone in colon carcinoma and toxicology research, not only for their relevance to human gastrointestinal studies but also for their role in providing detailed insights into the biliary pathway, the metabolism of xenobiotics within the colon, cancer and toxicology research.

**Organism** Human

**Tissue** Colon

**Disease** Adenocarcinoma

**Applications** Model of the GI (gastrointestinal) tract, measurement of the Trans-Epithelial/Endothelial Electrical Resistance (TEER). Caco-2 cells develop high TEER values of up to 2000 cm<sup>2</sup> (as measured by CLS using the CellZscope, nanoAnalytics, Münster, Germany).

**Synonyms** CaCo-2, CACO-2, Caco 2, CACO 2, CACO2, CaCo2, CaCO2, Caco2, Caco-II

### Characteristics

**Age** 72 years

**Gender** Male

**Ethnicity** Caucasian

## Caco-2 Cells | 300137

**Morphology** Epithelial-like

**Growth properties** Adherent

### Regulatory Data

**Citation** CaCo-2 (Cytion catalog number 300137)

**Biosafety level** 1

**NCBI\_TaxID** 9606

**CellosaurusAccession** CVCL\_0025

### Biomolecular Data

**Receptors expressed** Heat stable enterotoxin (Sta, E. coli), epidermal growth factor (EGF), retinoic acid binding protein I and retinol binding protein II, keratin positive.

**Antigen expression** Blood Type O, Rh+, HLA class II negative

**Isoenzymes** Me-2, 1, PGM3, 1, PGM1, 1, ES-D, 1, AK-1, 1, GLO-1, 1, G6PD, B.

**Tumorigenic** Yes, in nude mice. Form moderately well differentiated adenocarcinomas consistent with colonic primary (grade II)

**Virus resistance** Human immunodeficiency virus (HIV, LAV)

**Ploidy status** (P14), hypertetraploid

**MSI-status** Stable (MSS)

### Handling

**Culture Medium** EMEM (MEM Eagle), w: 2 mM L-Glutamine, w: 2.2 g/L NaHCO<sub>3</sub>, w: EBSS (Cytion article number 820100a)

**Supplements** Supplement the medium with 10% FBS and 1% NEAA

## Caco-2 Cells | 300137

**Dissociation Reagent**      Accutase

**Doubling time**      60 to 70 hours

**Subculturing**      Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

**Seeding density**       $1 \times 10^4$  cells/cm<sup>2</sup> will result in a 90% confluent monolayer in about 4 days

**Post-Thaw Recovery**      After thawing, plate the cells at  $5 \times 10^4$  cells/cm<sup>2</sup> and allow the cells to recover from the freezing process and to adhere for at least 24 hours.

**Freeze medium**      As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

## Caco-2 Cells | 300137

### Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below  $-150^{\circ}\text{C}$  to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a  $37^{\circ}\text{C}$  water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at  $300 \times g$  for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

### Incubation Atmosphere

$37^{\circ}\text{C}$ , 5%  $\text{CO}_2$ , humidified atmosphere.

### Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately  $-78^{\circ}\text{C}$  throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

### Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about  $-150$  to  $-196^{\circ}\text{C}$ . Storage at  $-80^{\circ}\text{C}$  is acceptable only as a short interim step before transfer to liquid nitrogen.

## Quality Control & Molecular Analysis

## Caco-2 Cells | 300137

### **Sterility**

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.