

**HROC24 T1 M1 Cells | 300813****General information**

<b>Description</b>	This is one of a series of tumor cell lines which have been established by PD Dr. Michael Linnebacher from PDTx (Patient-derived Tumor xenografts) since 2006.
<b>Organism</b>	Human
<b>Tissue</b>	Colon, Established from a PDx (patient-derived xenograft) of primary CRC tissue (Colon ascendens, UICC I, TNM stage T2N0M0R0L0V1 grading G2). Lk(n) + 0, $\Sigma$ Lk(n) 13
<b>Disease</b>	Epithelial
<b>Synonyms</b>	HROC24x

**Characteristics**

<b>Age</b>	98 years
<b>Gender</b>	Male
<b>Ethnicity</b>	Caucasian
<b>Morphology</b>	Small cells in colonies
<b>Growth properties</b>	Adherent

**Regulatory Data**

<b>Citation</b>	HROC24 T1 M1 (Cytion catalog number 300813)
<b>Biosafety level</b>	1
<b>NCBI_TaxID</b>	9606
<b>CellosaurusAccession</b>	CVCL_1U80

**Biomolecular Data**

<b>Tumorigenic</b>	Yes, in immune-suppressed nude mice
--------------------	-------------------------------------

**HROC24 T1 M1 Cells | 300813**

**Viruses** Free of human pathogenic viruses SV40, JC/BK, HBV, HCV, HIV.

**Ploidy status** Euploid

**MSI-status** MSI-H

**Mutational profile** APCmut, p53wt, K-Raswt, N-Raswt, H-Ras wt , B-RAFV600E, PIK3CAwt

**Handling**

**Culture Medium** DMEM:Ham's F12 (1:1), w: 3.1 g/L Glucose, w: 2.5 mM L-Glutamine, w: 15 mM HEPES, w: 0.5 mM Sodium pyruvate, w: 1.2 g/L NaHCO<sub>3</sub> (Cytion article number 820400a)

**Supplements** Supplement the medium with 10% FBS

**Dissociation Reagent** Accutase

**Doubling time** 20 hours

**Subculturing** Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

**Seeding density**  $2 \times 10^4$  cells/cm<sup>2</sup>

**Fluid renewal** Every 3 to 5 days

**Post-Thaw Recovery** Very fast

**Freeze medium** As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

## HROC24 T1 M1 Cells | 300813

### Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below  $-150^{\circ}\text{C}$  to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a  $37^{\circ}\text{C}$  water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at  $300 \times g$  for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

### Incubation Atmosphere

$37^{\circ}\text{C}$ , 5%  $\text{CO}_2$ , humidified atmosphere.

### Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately  $-78^{\circ}\text{C}$  throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

### Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about  $-150$  to  $-196^{\circ}\text{C}$ . Storage at  $-80^{\circ}\text{C}$  is acceptable only as a short interim step before transfer to liquid nitrogen.

## Quality Control & Molecular Analysis

## HROC24 T1 M1 Cells | 300813

### **Sterility**

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.