

A498 Cells | 300113

General information

Description

A498 cells are a human renal cell carcinoma cell line derived from the kidney tissue of a 58-year-old Caucasian male. These cells are extensively used in research related to kidney cancer, particularly for studying clear cell renal cell carcinoma, which is the most common type of kidney cancer in adults.

The A498 cell line is characterized by its epithelial-like morphology and has been a valuable model for investigating the molecular and cellular mechanisms of renal carcinogenesis. These cells exhibit several features typical of kidney cancer, including alterations in the expression of genes involved in cell cycle regulation, apoptosis, and angiogenesis.

A498 cells are particularly useful for examining the metabolic pathways altered in kidney cancer, as they display a distinct metabolic profile that includes changes in lipid and glucose metabolism. This aspect makes them suitable for metabolic targeting studies, which explore how altering metabolic pathways can inhibit tumor growth.

Furthermore, A498 cells are employed in drug discovery and toxicology studies to test the efficacy of new chemotherapeutic agents and targeted therapies. They are also used to study the response of renal cancer cells to hypoxic conditions—a common feature of solid tumors that significantly influences tumor behavior and treatment response.

Overall, the A498 cell line serves as an essential tool in renal cancer research, facilitating the development of more effective therapeutic strategies and enhancing our understanding of kidney cancer biology.

Organism Human

Tissue Kidney

Disease Renal cell carcinoma

Synonyms A-498

Characteristics

Age 52 years

Gender Male

Ethnicity Caucasian

Morphology Epithelial-like

Growth properties Monolayer, adherent

A498 Cells | 300113

Regulatory Data

Citation	A498 (Cytion catalog number 300113)
Biosafety level	1
NCBI_TaxID	9606
CellosaurusAccession	CVCL_1056

Biomolecular Data

Isoenzymes	PGM3, 1, PGM1, 1-2, ES-D, 2, Me-2, 1, AK-1, 1, GLO-1, 2, G6PD, B
Tumorigenic	Yes, in nude mice. Forms undifferentiated carcinoma, also forms tumors in anti thymocyte serum treated newborn mice
Ploidy status	Bimodal, tetraploid
MSI-status	Stable (MSS)

Handling

Culture Medium	EMEM (MEM Eagle), w: 2 mM L-Glutamine, w: 2.2 g/L NaHCO ₃ , w: EBSS (Cytion article number 820100a)
Supplements	Supplement the medium with 10% FBS and 1% NEAA
Dissociation Reagent	Accutase
Doubling time	62 hours
Subculturing	Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.
Seeding density	1 x 10 ⁴ cells/cm ² will result in a confluent monolayer within 4 days.

A498 Cells | 300113

Fluid renewal

Every 3 days

Post-Thaw Recovery

After thawing, plate the cells at 2×10^4 cells/cm² and allow the cells to recover from the freezing process and to adhere for at least 24 to 48 hours.

Freeze medium

As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C, 5% CO₂, humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

A498 Cells | 300113

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196 °C. Storage at -80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.