

MALME-3M Cells | 305583

Informações gerais

Description

The MALME-3M cell line is a human melanoma model used extensively in cancer research to investigate mechanisms of melanoma progression, immune evasion, and drug resistance. This cell line is derived from a metastatic melanoma lesion and demonstrates several characteristics relevant to aggressive melanoma, including its ability to express key oncogenic markers such as HER2 and its role in modulating the tumor microenvironment. Studies involving MALME-3M have highlighted its responsiveness to targeted therapies, such as bispecific antibodies aimed at HER2, and its use in evaluating T-cell mediated immunotherapies.

One significant area of research involving MALME-3M cells is their utility in studying the mechanisms of immune evasion in melanoma. For instance, co-culture systems pairing MALME-3M with immune cells allow researchers to explore how melanoma cells modulate immune responses through pathways like PD-1/PD-L1 and other immune checkpoint inhibitors. This cell line has also been genetically modified to study the effects of gene perturbations on immune interactions, making it a valuable tool for high-throughput genetic screening.

In addition to its role in immunological studies, MALME-3M cells are instrumental in exploring the effects of growth hormone (GH) on melanoma progression. Research has demonstrated that GH can enhance drug resistance and metastatic potential in MALME-3M cells by altering the composition of melanoma-derived exosomes. These exosomes can transfer drug resistance and migration-promoting factors to other cells in the tumor microenvironment. Such studies underscore the potential of targeting GH signaling pathways as a therapeutic strategy for overcoming melanoma chemoresistance.

Organism Human

Tissue Skin

Disease Melanoma

Metastatic site Lung

Synonyms Malme-3M, MALME 3M, Malme-3 M, MALME.3M, Malme3M, MALME3M, Malme-3 Monolayer

Características

Age 43 years

Gender Male

Ethnicity Caucasian

Morphology Fibroblast-like

Cell type Fibroblast

MALME-3M Cells | 305583

Growth properties Adherent

Dados regulatórios

Citation MALME-3M (Cytion catalog number 305583)

Biosafety level 1

NCBI_TaxID 9606

CellosaurusAccession CVCL_1438

Dados biomoleculares

Antigen expression HLA A2, Aw30, B13, B40(+/-), DRw7

Tumorigenic Yes, in nude mice

Manuseio

Culture Medium IMDM, w: 4.5 g/L Glucose, w: 4 mM L-Glutamine, w: 25 mM HEPES, w: 1.0 mM Sodium pyruvate, w: 3.024 g/L NaHCO₃ (Cytion article number 820800a)

Supplements Supplement the medium with 20% FBS

Dissociation Reagent Accutase

Subculturing Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with TrypLE Express, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

Seeding density 3×10^4 cells/cm²

Fluid renewal 2 to 3 times per week

MALME-3M Cells | 305583

Freeze medium

As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $300 \times g$ for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C , 5% CO_2 , humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

MALME-3M Cells | 305583

Controle de Qualidade e Análise Molecular

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.