

NCI-H1703 Cells | 305090

Informações gerais

Description

NCI-H1703 is a human lung cancer cell line derived from a stage I lung squamous cell carcinoma resected from a 54-year-old male smoker. This line exhibits a squamous epithelial morphology characteristic of lung carcinoma, reflecting its origin in lung tumor tissue:contentReference. It is classified as a non-small cell lung cancer (NSCLC) model, providing researchers with an in vitro representation of the squamous subtype of lung cancer.

NCI-H1703 is widely used in lung cancer research, particularly to study the squamous cell subtype of NSCLC:contentReference. It serves as a valuable model for investigating tumor biology and key cellular signaling pathways involved in lung cancer progression:contentReference. Researchers also employ NCI-H1703 to evaluate the efficacy of candidate anti-cancer compounds, including various targeted therapies. Additionally, this cell line can be used to establish xenograft models for preclinical testing of new treatments and to examine mechanisms of drug response and resistance in lung cancer.

Organism

Human

Tissue

Lung

Disease

Lung squamous cell carcinoma

Synonyms

NCI-H1703, H-1703, NCIH1703

Características

Age

54 years

Gender

Male

Ethnicity

European

Morphology

Epithelial

Growth properties

Adherent

Dados regulatórios

Citation

NCI-H1703 (Cytion catalog number 305090)

Biosafety level

1

NCI-H1703 Cells | 305090

NCBI_TaxID	9606
------------	------

CellosaurusAccession	CVCL_1490
----------------------	-----------

Dados biomoleculares

Manuseio

Culture Medium	RPMI 1640, w: 2.0 mM stable Glutamine, w: 2.0 g/L NaHCO ₃ (Cytion article number 820700a)
-----------------------	--

Supplements	Supplement the medium with 10% FBS
--------------------	------------------------------------

Dissociation Reagent	Accutase
-----------------------------	----------

Subculturing	Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.
---------------------	---

Fluid renewal	2 to 3 times per week
----------------------	-----------------------

Freeze medium	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.
----------------------	---

NCI-H1703 Cells | 305090

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $300 \times g$ for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C , 5% CO_2 , humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

Controle de Qualidade e Análise Molecular

NCI-H1703 Cells | 305090

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.