

## SNU-1 Cells | 305076

## Informações gerais

**Description**

The SNU-1 cell line is derived from the gastric carcinoma of a human adult and is widely utilized in gastric cancer research. This cell line provides an important model for studying the molecular and cellular mechanisms underlying gastric adenocarcinoma, a common and often deadly form of stomach cancer. SNU-1 cells are particularly valuable for investigating the genetic alterations and signaling pathways involved in the pathogenesis of gastric cancer, as well as for developing and testing new therapeutic strategies.

SNU-1 cells exhibit an epithelial morphology and are characterized by the expression of markers typical of gastric epithelial cells and adenocarcinoma, such as carcinoembryonic antigen (CEA) and cytokeratins. They are often used in studies exploring the role of oncogenes, tumor suppressor genes, and other molecular factors in gastric cancer progression. Researchers employ SNU-1 cells to assess the efficacy and mechanisms of action of chemotherapeutic agents, targeted therapies, and combination treatments. Additionally, SNU-1 cells serve as a model for understanding the tumor microenvironment and the interactions between cancer cells and stromal cells. The relevance of the SNU-1 cell line in gastric cancer research highlights its importance in advancing our knowledge of this malignancy and in the development of effective treatments for gastric cancer patients.

**Organism**

Human

**Tissue**

Stomach

**Disease**

Adenocarcinoma

**Synonyms**

SNU1, NCI-SNU-1

## Características

**Age**

44 years

**Gender**

Male

**Ethnicity**

Asian

**Morphology**

Epithelial

**Growth properties**

Suspension

## Dados regulatórios

**Citation**

SNU-1 (Cytion catalog number 305076)

## SNU-1 Cells | 305076

**Biosafety level** 1**NCBI\_TaxID** 9606**CellosaurusAccession** CVCL\_0099**Dados biomoleculares****Receptors expressed** Vasoactive intestinal peptide(VIP), expressed**Antigen expression** Blood Type O, Rh -, The cells express the surface glycoproteins carcinoembryonic antigen(CEA) and TAG 72.**Manuseio****Culture Medium** RPMI 1640, w: 2.0 mM stable Glutamine, w: 2.0 g/L NaHCO<sub>3</sub> (Cytion article number 820700a)**Supplements** Supplement the medium with 10% heat-inactivated FBS**Dissociation Reagent** Accutase**Subculturing** Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.**Seeding density** 0,3-1 x 10<sup>6</sup> cells/ml**Fluid renewal** 2 to 3 times per week**Post-Thaw Recovery** After thawing, plate the cells at 5 x 10<sup>4</sup> cells/cm<sup>2</sup> and allow the cells to recover from the freezing process and to adhere for at least 24 hours.**Freeze medium** As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

## SNU-1 Cells | 305076

### Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below  $-150^{\circ}\text{C}$  to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a  $37^{\circ}\text{C}$  water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at  $300 \times g$  for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

### Incubation Atmosphere

$37^{\circ}\text{C}$ , 5%  $\text{CO}_2$ , humidified atmosphere.

### Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately  $-78^{\circ}\text{C}$  throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

### Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about  $-150$  to  $-196^{\circ}\text{C}$ . Storage at  $-80^{\circ}\text{C}$  is acceptable only as a short interim step before transfer to liquid nitrogen.

## Controle de Qualidade e Análise Molecular

**SNU-1 Cells | 305076**

**Sterility**

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.