

## C-33 A Cells | 305072

## Informações gerais

## Description

C-33 A cells originate from the cervical tissue of a 66-year-old Caucasian female diagnosed with uterine cancer. This cell line is characterized by a unique genetic alteration in the TP53 gene, where a point mutation at codon 273 results in an arginine to cysteine substitution, leading to elevated expression of the p53 protein. This mutation plays a critical role in the pathophysiology of the cells, influencing their growth properties and tumorigenic potential.

Notably, C-33 A cells are confirmed to be tumorigenic. When introduced into immunodeficient nude mice, these cells have the capability to form undifferentiated carcinomas, which highlights their utility in cancer research, particularly in studies aimed at understanding the mechanisms of tumor initiation and progression in cervical cancer. Furthermore, these cells are negative for both DNA and RNA of human papillomavirus (HPV), distinguishing them from many other cervical cancer cell lines which often carry HPV integrations. This aspect makes C-33 A cells particularly valuable for studying cervical cancer that develops independently of HPV infection, offering insights into alternative pathways of carcinogenesis.

## Organism

Human

## Tissue

Cervix

## Disease

Squamous cell carcinoma of the cervix uteri

## Synonyms

C33A, C33a, C33-A, C-33-A, C-33A, C33

## Características

## Age

66 years

## Gender

Female

## Morphology

Epithelial

## Growth properties

Adherent

## Dados regulatórios

## Citation

C33A (Cytion catalog number 305072)

## Biosafety level

1

## NCBI\_TaxID

9606

## C-33 A Cells | 305072

CellosaurusAccession CVCL\_1094

## Dados biomoleculares

**Protein expression** Oncogenes: P53 , Prb**Tumorigenic** Yes

## Manuseio

**Culture Medium** EMEM (MEM Eagle), w: 2 mM L-Glutamine, w: 2.2 g/L NaHCO<sub>3</sub>, w: EBSS (Cytion article number 820100a)**Supplements** Supplement the medium with 10% FBS and 1% NEAA**Dissociation Reagent** Accutase**Subculturing** Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.**Fluid renewal** 2 to 3 times per week**Freeze medium** As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

**C-33 A Cells | 305072**

**Thawing and  
Culturing Cells**

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

**Incubation  
Atmosphere**

37°C, 5% CO<sub>2</sub>, humidified atmosphere.

**Shipping  
Conditions**

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

**Storage  
Conditions**

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196 °C. Storage at -80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

**Controle de Qualidade e Análise Molecular**

**C-33 A Cells | 305072**

**Sterility**

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.