

SNU-668 Cells | 305635

Description

The SNU-668 cell line is a human gastric carcinoma model originally derived from the poorly differentiated adenocarcinoma tissue of the stomach. This cell line has been widely used in studies of gastric cancer pathogenesis, signaling mechanisms, and drug responsiveness. Genomic characterization reveals that SNU-668 carries frequent mutations and chromosomal aberrations commonly observed in diffuse-type gastric cancers. Notably, it shows alterations in key oncogenic pathways such as TP53 mutation and possible activation of PI3K/AKT signaling, which may contribute to its tumorigenic properties and therapy resistance.

SNU-668 has also been included in comprehensive multi-omics profiling projects such as the Cancer Cell Line Encyclopedia (CCLE), where it was assessed for transcriptomic, genomic, methylation, and proteomic signatures. The cell line exhibits distinct DNA methylation patterns and global histone modification profiles, which may play roles in epigenetic regulation of gene expression. Additionally, analysis of dependency maps has suggested lineage-specific vulnerabilities that could inform targeted therapy strategies for diffuse gastric carcinomas. As a model for stomach cancer with Asian ethnic background origin, SNU-668 continues to be an important tool in the preclinical evaluation of molecularly guided therapeutics.

Organism

Human

Tissue

Gastric

Disease

signet ring cell adenocarcinoma

Metastatic site

Ascites

Synonyms

SNU668, NCI-SNU-668

Age

63 years

Gender

Male

Ethnicity

Korean

Morphology

Epithelial-like

Cell type

Epithelial

Growth properties

Adherent, monolayer

SNU-668 Cells | 305635

Citation SNU-668 (Cytion catalog number 305635)

Biosafety level 1

NCBI_TaxID 9606

CellosaurusAccession CVCL_5081

Mutational profile Mutation: KRAS, Simple, p.Gln61Lys (c.181C>A), Homozygous; Mutation: TP53, Simple, p.Ser215Asn (c.644G>A), Homozygous

Culture Medium RPMI 1640, w: 2.0 mM stable Glutamine, w: 2.0 g/L NaHCO₃ (Cytion article number 820700a)

Supplements Supplement the medium with 10% heat inactivated FBS

Dissociation Reagent Accutase

Doubling time 26 hours

Subculturing Remove medium, add fresh 0.25 % trypsin 0.02 % EDTA solution, stand culture flask at 37°C for 3 to 5 minutes, add culture medium and collect the cells, transfer the medium into 15ml tube, centrifuge, aspirate the medium, resuspend the pellets with culture medium and dispense into the culture flask

Fluid renewal 2 to 3 times per week

Freeze medium As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

SNU-668 Cells | 305635

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $300 \times g$ for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C , 5% CO_2 , humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

SNU-668 Cells | 305635

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.