

## HT-29 MTX E12 Cells | 305801

### Description

HT-29-MTX-E12 is a goblet cell-like subclone derived from the human colorectal adenocarcinoma cell line HT29 through selection with methotrexate (MTX), a process that induces differentiation toward mucus-secreting phenotypes. Among several subclones developed from HT29-MTX, the E12 subclone stands out due to its robust formation of confluent monolayers with tight junctions and a significantly thick, continuous mucus layer on the apical surface. This subclone features a higher proportion of mature goblet cells, as demonstrated by Alcian Blue staining, transmission electron microscopy (TEM), and expression of mucin genes MUC1 and MUC2. In fact, MUC1 and MUC2 mRNA levels were substantially higher in HT-29-MTX-E12 compared to other subclones and parent HT29 cells, correlating with a mucus thickness of approximately  $142 \pm 51 \mu\text{m}$ -comparable to the in vivo intestinal environment.

Functionally, HT-29-MTX-E12 has been shown to model the barrier properties of the human intestinal mucus layer, particularly in evaluating the absorption of lipophilic drugs. The presence of a thick mucus barrier significantly reduces the apparent permeability coefficients (Papp) of lipophilic compounds such as testosterone and various barbiturates when compared to mucus-free Caco-2 cells. For example, testosterone showed a 43% reduction in Papp in HT-29-MTX-E12, highlighting the impact of mucus on drug diffusion. Despite having a leakier epithelial barrier than Caco-2 cells, HT-29-MTX-E12 maintains physiological relevance through its mucus-producing capacity, making it a valuable in vitro model for investigating intestinal drug absorption and the influence of mucus on permeability.

**Organism** Human

**Tissue** Colon

**Disease** Colon adenocarcinoma

**Synonyms** HT29-MTX-E12, MTX-E12

**Age** 44 years

**Gender** Female

**Ethnicity** Caucasian

**Cell type** Epithelial

**Growth properties** Adherent

## HT-29 MTX E12 Cells | 305801

<b>Citation</b>	HT-29-MTX-E12 (Cytion catalog number 305801)
<b>Biosafety level</b>	1
<b>NCBI_TaxID</b>	9606
<b>CellosaurusAccession</b>	CVCL_G356
<b>Mutational profile</b>	Mutation: APC, Simple, p.Glu853Ter (c.2557G>T), Heterozygous (from parent cell line).Mutation, APC, Simple, p.Thr1556Asnfs*3 (c.4666dupA) (c.4666_4667insA), Heterozygous (from parent cell line).Mutation, BRAF, Simple, p.Val600Glu (c.1799T>A), Heterozygous (from parent cell line).Mutation, PIK3CA, Simple, p.Pro449Thr (c.1345C>A), Heterozygous (from parent cell line).Mutation, SMAD4, Simple, p.Gln311Ter (c.931C>T), Homozygous (from parent cell line).Mutation, TP53, Simple, p.Arg273His (c.818G>A), Homozygous (from parent cell line).
<b>Culture Medium</b>	EMEM (MEM Eagle), w: 2 mM L-Glutamine, w: 2.2 g/L NaHCO <sub>3</sub> , w: EBSS (Cytion article number 820100a)
<b>Supplements</b>	Supplement the medium with 10% FBS and 1% NEAA
<b>Dissociation Reagent</b>	Accutase
<b>Freeze medium</b>	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

## HT-29 MTX E12 Cells | 305801

### Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below  $-150^{\circ}\text{C}$  to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a  $37^{\circ}\text{C}$  water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at  $300 \times g$  for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

### Incubation Atmosphere

$37^{\circ}\text{C}$ , 5%  $\text{CO}_2$ , humidified atmosphere.

### Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately  $-78^{\circ}\text{C}$  throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

### Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about  $-150$  to  $-196^{\circ}\text{C}$ . Storage at  $-80^{\circ}\text{C}$  is acceptable only as a short interim step before transfer to liquid nitrogen.

## HT-29 MTX E12 Cells | 305801

### **Sterility**

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.