

CHO-STEAP1 Cells | 305983

Description

Disclaimer: The prices displayed for cell lines are exclusively for academic/not-for-profit customers. For commercial entities the price is approximately €6,250.

If you represent a commercial entity or are unsure which category applies, please [contact us](#).

CHO-STEAP1 cells are recombinant Chinese hamster ovary (CHO) cells engineered to stably express human six-transmembrane epithelial antigen of the prostate 1 (STEAP1), a cell surface protein highly associated with multiple solid tumors. STEAP1 is a member of the STEAP family of metalloredoxases and is characterized by six transmembrane domains with localization predominantly at the plasma membrane and intracellular vesicular compartments. Although its precise physiological function remains incompletely understood, STEAP1 has been implicated in intercellular communication, metal ion homeostasis, redox regulation, and tumor cell proliferation. Elevated STEAP1 expression has been reported in prostate cancer, Ewing sarcoma, bladder cancer, lung cancer, and several other malignancies, making it an important target in oncology-focused therapeutic development.

CHO-STEAP1 cells are widely used for development and characterization of STEAP1-targeted therapeutics, including monoclonal antibodies, antibody-drug conjugates, bispecific T-cell engagers, radioligand therapies, and engineered immune cell approaches such as CAR-T and CAR-NK therapies. The stable recombinant expression system enables quantitative analysis of antibody binding affinity, receptor occupancy, antigen density, internalization behavior, and target-specific cytotoxicity. These cells are also valuable for flow cytometry assay development, epitope mapping, high-throughput screening, and validation of STEAP1-targeted imaging agents. Because CHO cells provide a robust and relatively low-background platform for recombinant protein expression, CHO-STEAP1 models are frequently used for standardized assay development and preclinical therapeutic evaluation.

Organism Chinese hamster

Tissue Ovary

Disease Chinese hamster ovary, non-neoplastic; genetically engineered for STEAP1 surface expression

Applications Antibody screening; STEAP1-targeted therapy development; ADC development; prostate/bladder cancer research; flow cytometry

Age Adult

Gender Female

Morphology Epithelial-like

CHO-STEAP1 Cells | 305983

Cell type	Epithelial cells
Growth properties	Adherent/suspension
Citation	CHO-STEAP1 (Cytion catalog number 305983)
Biosafety level	1
NCBI_TaxID	10029
CellosaurusAccession	CVCL_A8X2
GMO Status	GMO-S1: This CHO cell line contains a STEAP1 expression cassette supporting receptor-function analyses. This classification applies only within Germany and may differ elsewhere.
Receptors expressed	STEAP1
Culture Medium	<p>For adherent cultures: DMEM:Ham's F12 (1:1), w: 3.1 g/L Glucose, w: 2.5 mM L-Glutamine, w: 15 mM HEPES, w: 0.5 mM Sodium pyruvate, w: 1.2 g/L NaHCO₃ (Cytion article number 820400a)</p> <p>For suspension cultures: CHO Growth Medium A (from InSCREENeX; InSCREENeX catalog number INS-ME-1039)</p>
Supplements	For adherent cultures: Supplement the medium with 5% FBS. Add Geneticin (G418-Sulfat) to achieve a final concentration of 0.5 mg/mL.
Dissociation Reagent	For adherent cultures: Trypsin-EDTA
Doubling time	approx. 14-16 hours

CHO-STEAP1 Cells | 305983

Subculturing For routine adherent cell culture: Aspirate the old culture medium from the adherent cells, and wash them with PBS to remove any remaining medium. After aspirating the PBS, add the appropriate volume of Trypsin/EDTA solution based on the culture vessel size (e.g., 1 ml for a T25 flask, 3 ml for a T75 flask) and incubate at room temperature or 37°C for 5-10 minutes, or until the cells detach. Monitor detachment under a microscope, and gently tap the vessel if necessary to release the cells. Once detached, add complete medium to inactivate the Trypsin/EDTA, gently resuspend the cells, and transfer an aliquot of the cell suspension into a new culture vessel containing fresh medium. Place the vessel in an incubator set to 37°C with 5% CO₂, and change the medium every 2-3 days.

Split ratio 1 to 5

Seeding density 2 to 5 x 10⁴ cells/cm²

Fluid renewal 2 to 3 times per week

Post-Thaw Recovery After thawing, split the cells at a ratio of 1:2 to 1:3 in T25 flasks and allow the cells to recover from the freezing process and to adhere (for adherent cultures) for at least 24 hours.

Freeze medium As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

CHO-STEAP1 Cells | 305983

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $300 \times g$ for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C , 5% CO_2 , humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

CHO-STEAP1 Cells | 305983

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.