

Kelly Cells | 300317

Description

The Kelly cell line is a human neuroblastoma cell line derived from a tumor biopsy. Neuroblastoma is a malignant tumor that arises from neural crest cells, typically affecting children and infants. Kelly cells are utilized extensively in research due to their aggressive growth characteristics and their ability to differentiate into neuron-like cells under specific conditions. These cells exhibit properties typical of neuroblastoma, including high levels of MYCN amplification, which is associated with poor prognosis and aggressive tumor behavior. This makes the Kelly cell line a valuable model for studying the molecular mechanisms of neuroblastoma and for testing potential therapeutic agents.

Kelly cells are adherent in culture and can grow in a monolayer, making them suitable for a wide range of experimental applications, including drug screening, gene expression studies, and investigations into cellular signaling pathways. They are particularly useful for studying the effects of MYCN-driven oncogenesis and for evaluating the efficacy of targeted therapies against neuroblastoma. The Kelly cell line also serves as a model for understanding the biology of neuroblastoma metastasis, as these cells have the capability to migrate and invade, reflecting the behavior of aggressive neuroblastoma in vivo.

Organism Human

Tissue Brain

Disease Neuroblastoma

Synonyms KELLY, NB19, NB-19, NB19-RIKEN

Age 1 year

Gender Female

Ethnicity Caucasian

Growth properties Adherent

Citation Kelly (Cytion catalog number 300317)

Biosafety level 1

Kelly Cells | 300317

NCBI_TaxID	9606
CellosaurusAccession	CVCL_2092
Tumorigenic	Yes, in nude mice.
Viruses	Negative for HPV (Human Papilloma Virus)
Products	N-myc RnA
Culture Medium	RPMI 1640, w: 2.0 mM stable Glutamine, w: 2.0 g/L NaHCO ₃ (Cytion article number 820700a)
Supplements	Supplement the medium with 10% FBS
Dissociation Reagent	Accutase
Doubling time	30 hours
Subculturing	Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.
Seeding density	1 x 10 ⁴ cells/cm ²
Fluid renewal	2 to 3 times per week
Freeze medium	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

Kelly Cells | 300317

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $300 \times g$ for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C , 5% CO_2 , humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

Kelly Cells | 300317

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.