

MRC-9 Cells | 300394

Description

The MRC-9 cell line is a human diploid fibroblast strain derived from the lungs of a White female fetus at 15 weeks gestation. It was established for long-term use in biological research, vaccine production, and diagnostic virology. The cells exhibit adherent growth properties and maintain a stable, diploid karyotype throughout serial propagation. Importantly, MRC-9 cells possess normal chromosomal features and are non-tumorigenic under typical conditions. However, they have demonstrated tumorigenic properties when tested in semisolid media, but not when implanted in immunosuppressed mice.

MRC-9 cells have been widely used for viral susceptibility studies, showing broad susceptibility to various viruses such as poliovirus, adenovirus, rubella, and cytomegalovirus. These cells are particularly valuable for their use in virology research and vaccine development. While their proliferation rate is slower compared to other human fibroblast strains like MRC-5, MRC-9 cells can still undergo a significant number of population doublings, with reports of up to 46 doublings.

Organism Human

Tissue Fetal lung

Synonyms MRC9, Medical Research Council cell strain-9

Age Fetus

Gender Female

Cell type Fibroblast

Growth properties Adherent

Citation MRC-9 (Cytion catalog number 300394)

Biosafety level 1

NCBI_TaxID 9606

CellosaurusAccession CVCL_2629

MRC-9 Cells | 300394

Culture Medium

EMEM (MEM Eagle), w: 2 mM L-Glutamine, w: 2.2 g/L NaHCO₃, w: EBSS (Cytion article number 820100a)

Supplements

Supplement the medium with 10% FBS and 1% NEAA

Dissociation Reagent

Accutase

Subculturing

Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

Freeze medium

As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

MRC-9 Cells | 300394

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $300 \times g$ for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C , 5% CO_2 , humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

MRC-9 Cells | 300394

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.