

AH-130 | 500412

Description Yoshida 1956 AH-130

Organism

Tissue

Disease

Metastatic site

Applications -

Synonyms Yoshida AH-130 Yoshida AH130 AH130 AH 130 AH-130 Yoshida AH130-TC AH130/P

Breed/Subspecies -

Age

Gender

Ethnicity Q -

Morphology

Cell type

Growth properties

Citation AH-130 Cytion 500412

Biosafety level 1

NCBI_TaxID 10116

AH-130 | 500412

CellosaurusAccession CVCL_4367

GMO Status 1956

Tumorigenic Wistar

Viruses RAP .

Virus susceptibility

Culture Medium DMEM Ham's F12 1:1 3.1 g/L 2.5 mM L- 15 mM HEPES w: 0.5 mM w: 1.2 g/L NaHCO₃ Cytion

Supplements 10% FBS

Dissociation Reagent Accutase

Doubling time 18 24 BD=

Subculturing 15 PBS T25 3-5 T75 5-10 Accutase T25 1-2 T75

Split ratio 1 3

Seeding density $2 \times 10^4 / \text{cm}^2$

Fluid renewal 3 5

Post-Thaw Recovery

Freeze medium FBS + 10% DMSO CM-1 Cytion 800100

AH-130 | 500412

Thawing and Culturing Cells	1.				
	2.		-150°C		3
	3.		37°C	40-60	
	4.			70%	
	5.		8	15	
	6.	300 x g	3		
	7.	10		T25	T25
	8.				

Incubation Atmosphere

37°C 5% CO₂

Shipping Conditions					-78 °C
----------------------------	--	--	--	--	--------

Storage Conditions

-150 -196 °C

-80 °C

Sterility					
		PCR			