

T47D Cells | 300353

Description

The T47D cell line, originating from the pleural effusion of an infiltrating ductal carcinoma of the breast, has become a critical resource in breast cancer research. T-47D cells are unique in the realm of cancer research for their hormonal expression profile, particularly for carrying receptors for 17 beta estradiol, various other steroids, and calcitonin. Additionally, T47D cells express the WNT7B oncogene.

T47D cells are notable for their progesterone receptor expression not being regulated by estradiol, despite the hormone's abundance within the cells, setting them apart from MCF7 cells, which are widely recognized for their estrogen receptor positivity and are frequently used to explore estrogen's role in tumor proliferation and response to therapies.

The utility of the T47D cell line extends to the formation of xenografts in immunodeficient mice, which are valuable for drug testing, observing receptor status changes, and studying angiogenesis.

Furthermore, the T-47D cell line is a resource for cancer gene studies, providing insights into the genomic and proteomic landscape that drives breast cancer. By facilitating a deeper understanding of the proteomic and transcriptomic profiles of breast cancer, the t47d breast cancer cell line aids in the identification of new breast cancer cell phenotypes and the development of targeted therapies.

T47D cells have been instrumental in studying the effects of hormones like progesterone on breast cancer, offering insights into transcriptional regulation, drug resistance, and the development of xenograft models for therapeutic testing.

Organism Human

Tissue Breast

Disease Invasive ductal carcinoma

Metastatic site Pleural effusion

Synonyms T-47-D, T47-D, T47D:A, T47D

Age 54 years

Gender Female

Ethnicity Caucasian

Morphology Epithelial-like

T47D Cells | 300353

Growth properties Monolayer, adherent

Citation T47D (Cytion catalog number 300353)

Biosafety level 1

NCBI_TaxID 9606

CellosaurusAccession CVCL_0553

Receptors expressed Estradiol, steroids, calcitonin, androgen, progesterone, glucocorticoid, prolactin, estrogen

Isoenzymes G6PD, B, PGM1, 1, PGM3, 1, ES-D, 2, Ak-1, 1, GLO-1, 1-2

Oncogenes Wnt3 +, wnt7h +, wnt7b+

Tumorigenic Yes, in nude mice

Mutational profile TP53 mut

Karyotype Mode = 66, dicentric and extra long submetacentric chromosomes

Culture Medium RPMI 1640, w: 2.0 mM stable Glutamine, w: 2.0 g/L NaHCO₃ (Cytion article number 820700a)

Supplements Supplement the medium with 10% FBS, 10 microgram/ml HREC insulin

Dissociation Reagent Accutase

T47D Cells | 300353

Subculturing Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

Seeding density 1×10^4 cells/cm²

Fluid renewal 2 to 3 times per week

Post-Thaw Recovery After thawing, plate the cells at 5×10^4 cells/cm² and allow the cells to recover from the freezing process and to adhere for at least 24 hours.

Freeze medium As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

T47D Cells | 300353

**Thawing and
Culturing Cells**

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

**Incubation
Atmosphere**

37°C, 5% CO₂, humidified atmosphere.

**Shipping
Conditions**

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

**Storage
Conditions**

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196 °C. Storage at -80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

T47D Cells | 300353

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.