

A375 Cells | 300110

Description

The A375 human melanoma cell line, isolated from the skin of a 54-year-old female patient with malignant melanoma, is a significant resource in cancer research, particularly in the study of human melanoma, one of the most aggressive forms of skin cancer. The A375 cell line is known for its rapid growth rate and high tumorigenic potential, making them suitable for various experimental applications, including in vitro studies on cell proliferation, migration, and invasion, as well as in vivo tumorigenesis assays.

A375 cells exhibit high tumorigenic potential in immunosuppressed mice, forming rapidly growing amelanotic melanomas. The presence of the BRAFV600E mutation in A375 cells makes them highly sensitive to MEK inhibition, providing a valuable tool for investigating targeted therapies in melanoma treatment. The treatment of A375 cells with vemurafenib, for example, has been shown to enhance the induction of MHC Class I and Class II molecules, offering insights into the interactions between melanoma cells and the immune system.

In addition to their role in basic melanoma research, A375 cells are used in drug screening and in the investigation of signaling pathways involved in cancer cell survival, proliferation, and metastasis. A375 cells have further been utilized in apoptosis studies and A375 isogenic cell lines and the introduction of reporter proteins like-Luc (luc2) enable the study of gene function and the monitoring of cellular responses in real-time. A375 cells' suitability as a transfection host and their use in stable reporter cell lines also contribute to their versatility in research applications.

In summary, the A375 human melanoma cell line is a pivotal tool in the investigation of human melanoma, offering a comprehensive model for studying the molecular and cellular mechanisms underlying melanoma progression, the efficacy of therapeutic agents, and the interaction between cancer cells and the immune system.

Organism Human

Tissue Skin

Disease Melanoma

Synonyms A 375, A-375, A375-MEL, A375-mel, A375mel

Age 54 years

Gender Female

Morphology Epithelial-like

Growth properties Adherent

A375 Cells | 300110

Citation	A375 (Cytion catalog number 300110)
Biosafety level	1
NCBI_TaxID	9606
CellosaurusAccession	CVCL_0132
Antigen expression	P53 positive
Tumorigenic	Yes, in nude mice
Mutational profile	BRAF V600Emut
Karyotype	A375 cells are characterized by their hypotriploid karyotype, with a modal chromosome number of 62, and the presence of nine marker chromosomes in each cell, highlighting the genetic alterations associated with malignant melanoma.
Culture Medium	DMEM, w: 4.5 g/L Glucose, w: 4 mM L-Glutamine, w: 3.7 g/L NaHCO ₃ , w: 1.0 mM Sodium pyruvate (Cytion article number 820300a)
Supplements	Supplement the medium with 10% FBS
Dissociation Reagent	Accutase
Doubling time	20 hours
Subculturing	Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

A375 Cells | 300110

Seeding density 1×10^4 cells/cm² will result in a confluent monolayer within 4 days.

Fluid renewal 2 to 3 times per week

Post-Thaw Recovery After thawing, plate the cells at 4×10^4 cells/cm² and allow the cells to recover from the freezing process and to adhere for at least 24 hours.

Freeze medium As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere 37°C, 5% CO₂, humidified atmosphere.

A375 Cells | 300110

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.