Product sheet

CHO-TACD2 | 305415

Description	/ **6,250** CHO-TACD2 CHO TACD2 12,600 “ ” TACD2 TACD2 TROP2
Organism	
Tissue	
Disease	TACD2/TROP2 GA733-1
Applications	ADC TROP2 / /
Age	
Gender	
Morphology	
Cell type	
Growth properties	/
Citation	CHO-TACD2 Cytion 305415
Biosafety level	1
NCBI_TaxID	10029
CellosaurusAccession	CVCL_A8X3

Product sheet

CHO-TACD2 | 305415

GMO Status	GMO-S1 CHO TACD2
Receptors expressed	TACD2 TROP2 GA733-1
Culture Medium	DMEM:Ham's F12 (1:1) w: 3.1 g/L w: 2.5 mM L- w: 15 mM HEPES w: 0.5 mM w: 1.2 g/L NaHCO3 Cytio CHO A InSCREENeX InSCREENeX INS-ME-1039
Supplements	5% FBS G418- 0.5 mg/mL
Dissociation Reagent	-EDTA
Doubling time	14-16
Subculturing	PBS PB 5% 2-3 /EDTA T25 1 T75 3 37
Split ratio	1 5
Seeding density	2 5×10^{4} /
Fluid renewal	2 3
Post-Thaw Recovery	1:2 1:3 T25 24
Freeze medium	FBS +10% DMSO CM-1 Cytion 800100 CM-1 Cytion 800100

Thawing and Culturing Cells	1. 2. -150°C 3 3. 37°C 40-60 4. 70% 5. 8 15 6. 300 x g 3 7. 10 T25 T25 8.
Incubation Atmosphere	37°C, 5% $_{CO2}$,
Flask Coating	
Freezing Procedure	-78 °C
Shipping Conditions	-78 °C
Storage Conditions	-150 -196 -80 °C
Sterility	PCR