

CHO-CCR8 | 305418

Description	/ 6,250				
	CHO-CCR8	CHO	CCR8	8,000	CCR8
		CCR8			

Organism	
Tissue	

Disease	CCR8
----------------	------

Applications	CCR8
---------------------	------

Age	
------------	--

Gender	
---------------	--

Morphology	
-------------------	--

Cell type	
------------------	--

Growth properties	/
--------------------------	---

Citation	CHO-CCR8 Cytion 305418
-----------------	------------------------

Biosafety level	1
------------------------	---

NCBI_TaxID	10029
-------------------	-------

CellosaurusAccession	CVCL_A8V6
-----------------------------	-----------

CHO-CCR8 | 305418

GMO Status	GMO-S1	CHO	GPCR	CCR8
-------------------	--------	-----	------	------

Receptors expressed	CCR8	CHEMR1	CDw198
----------------------------	------	--------	--------

Culture Medium	DMEM:Ham's F12 (1:1)	w: 3.1 g/L	w: 2.5 mM L-	w: 15 mM HEPES	w: 0.5 mM	w: 1.2 g/L NaHCO3	Cytio
	CHO	A	InSCREENeX	InSCREENeX	INS-ME-1039		

Supplements	5% FBS	G418-	0.5 mg/mL
--------------------	--------	-------	-----------

Dissociation Reagent	-EDTA
-----------------------------	-------

Doubling time	14-16
----------------------	-------

Subculturing	PBS	PB5%	2-3	/EDTA	T25	1	T75	3	37
---------------------	-----	------	-----	-------	-----	---	-----	---	----

Split ratio	1	5
--------------------	---	---

Seeding density	2.5×10^4	/
------------------------	-------------------	---

Fluid renewal	2	3
----------------------	---	---

Post-Thaw Recovery	1:2	1:3	T25	24
---------------------------	-----	-----	-----	----

Freeze medium	FBS +10% DMSO	CM-1	Cytion	800100	CM-1	Cytion	800100
----------------------	---------------	------	--------	--------	------	--------	--------

CHO-CCR8 | 305418

Thawing and Culturing Cells				
1.				
2.		-150°C		3
3.		37°C	40-60	
4.			70%	
5.		8	15	
6.	300 x g	3		
7.	10		T25	T25
8.				

Incubation Atmosphere 37°C, 5% CO₂

Flask Coating

Freezing Procedure -78 °C

Shipping Conditions -78 °C

Storage Conditions -150 -196 -80 °C

/ /HLA

Sterility PCR