

**CHO-CXCR4 | 305411MH**

Description				
CHO-CXCR4-Medium-high	CHO	CXCR4	9500	
CXCR4 CD184		HIV	CXCL12	

Organism	
Tissue	

Disease	Chinese hamster ovary, non-neoplastic; genetically engineered for CXCR4 surface expression (medium-high expression level)
---------	---

Applications	Antibody screening; CXCR4-targeted therapy development; HIV entry research; hematopoietic stem cell biology; flow cytometry
--------------	---

Synonyms	CHO-CXCR4
----------	-----------

Age	
-----	--

Gender	
--------	--

Morphology	
------------	--

Cell type	Epithelial cells
-----------	------------------

Growth properties	/
-------------------	---

Citation	CHO-CXCR4 Medium-high Cytion 305411MH
----------	---------------------------------------

Biosafety level	1
-----------------	---

NCBI_TaxID	10029
------------	-------

**CHO-CXCR4 | 305411MH**

**CellosaurusAccession** CVCL\_A8W0

**GMO Status** GMO-S1: This CHO derivative contains a construct driving medium-to-high expression of human CXCR4 for GPCR signaling and ligand-binding analyses. This classification applies only within Germany and may differ elsewhere.

**Receptors expressed** CXCR4 CD184

**Culture Medium** A DMEM:Ham's F12 (1:1) w: 3.1 g/L w: 2.5 mM L- w: 15 mM HEPES w: 0.5 mM w: 1.2 g/L NaHCO3 Cytio  
 InSCREENeX InSCREENeX INS-ME-1039

**Supplements** 5% FBS G418- 0.5 mg/mL

**Dissociation Reagent** -EDTA

**Doubling time** approx. 14-16 hours

**Subculturing** PBS PBS /EDTA T25 1 T75 3 37

**Split ratio** 1 to 5

**Seeding density** 2 to 5 x 10<sup>4</sup> cells/cm<sup>2</sup>

**Fluid renewal** 2 3

**Post-Thaw Recovery** 1:2 1:3 T25 24

**Freeze medium** FBS +10% DMSO CM-1 Cytion 800100

CHO-CXCR4 | 305411MH

Thawing and Culturing Cells				
1.				
2.		-150°C		3
3.		37°C	40-60	
4.			70%	
5.		8	15	
6.	300 x g	3		
7.	10		T25	T25
8.				

**Incubation Atmosphere**

37°C, 5% CO<sub>2</sub>, humidified atmosphere.

**Shipping Conditions**

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

**Storage Conditions**

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196 °C. Storage at -80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

/ /HLA

**Sterility**

PCR