Product sheet

CHO-HER2 | 305413H

Description					
	CHO-HER2	CHO	HER2	85,000	HER2
Organism					
Tissue					
Disease	Chinese hamster ovary, non-neoplastic; genetically engineered for HER2 (ErbB2/CD340) surface expression (high expression level)				
Applications	Antibody screening; ADCC/CDC assays; HER2-targeted therapy development; breast/gastric cancer research; flow cytometry				
Synonyms	CHO-HER2				
Age					
Gender					
Morphology					
Cell type	Epithelial cells				
Growth properties	/				
Citation	CHO-HER2 High Cytion 305413H				
Biosafety level	1				
NCBI_TaxID	10029				

Product sheet

CHO-HER2 | 305413H

CellosaurusAccession	CVCL_A8W6
GMO Status	GMO-S1: This CHO cell line contains a construct enabling high-level expression of human HER2 for oncology and receptor-signaling studies. This classification applies only within Germany and may differ elsewhere.
Receptors expressed	HER2
Culture Medium	DMEM:Ham's F12 (1:1) w: 3.1 g/L w: 2.5 mM L- w: 15 mM HEPES w: 0.5 mM w: 1.2 g/L NaHCO3 Cytio A InSCREENeX InSCREENeX INS-ME-1039
Supplements	5% FBS G418- 0.5 mg/mL
Dissociation Reagent	-EDTA
Doubling time	approx. 14-16 hours
Subculturing	PBS PBS /EDTA T25 1 T75 3 37
Split ratio	1 to 5
Seeding density	2 to 5 x 10^4 cells/cm^2
Fluid renewal	2 3
Post-Thaw Recovery	1:2 1:3 T25 24
Freeze medium	FBS +10% DMSO CM-1 Cytion 800100

CHO-HER2 | 305413H

Thawing and Culturing Cells	1. 2. -150°C 3 3. 37°C 40-60 4. 70% 5. 8 15 6. 300 x g 3 7. 10 T25 T25 8.
Incubation Atmosphere	37°C, 5% CO_2, humidified atmosphere.
Shipping Conditions	Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately −78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.
Storage Conditions	For long-term preservation, place vials in vapor-phase liquid nitrogen at about −150 to −196 °C. Storage at −80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

/ /HLA

Sterility	PCR