

Product sheet

NCI-H2126 | 300639

Description	NCI-H2126	NSCLC			
	NCI-H2126		6q 13q	NSCLC	
	NCI-H2126				NSCLC

Organism

Tissue

Disease

Metastatic site

Applications

Synonyms H-2126 NCI-H2126 NCI-H2126

Age 65

Gender

Ethnicity

Morphology

Growth properties

Citation NCI-H2126 Cytion 300639

Biosafety level 2

NCBI_TaxID 9606

Product sheet

NCI-H2126 | 300639

CellosaurusAccession CVCL_1532

Isoenzymes AK-1 1 ES-D 1-2 G6PD B GLO-I 2 Me-2 0 PGM1 1-2 PGM3 2

Tumorigenic

Viruses EB

Ploidy status

Culture Medium DMEM:Ham's F12 1:1 w: 3.1 g/L w: 2.5 mM L- w: 15 mM HEPES w: 0.5 mM w: 1.2 g/L NaHCO3 Cytion 8

Supplements 5% FBS 0.005 mg/mL 0.01 mg/mL 30nM 10 nM 10 nM beta-estradiol

Dissociation Reagent Accutase

Subculturing PBS T25 3-5 PBS T75 5-10 Accutase T25 1-2 T75 2.5

Freeze medium FBS +10% DMSO CM-1 Cytion 800100 CM-1 Cytion 800100

Product sheet

NCI-H2126 | 300639

Thawing and
Culturing Cells

- | | | | | |
|----|---------|--------|-------|-----|
| 1. | | | | |
| 2. | | -150°C | | 3 |
| 3. | | 37°C | 40-60 | |
| 4. | | 70% | | |
| 5. | | 8 | 15 | |
| 6. | 300 x g | 3 | | |
| 7. | 10 | | T25 | T25 |
| 8. | | | | |

Incubation
Atmosphere

37°C, 5% CO₂

Flask Coating

Freezing
Procedure

-78 °C

Shipping
Conditions

-78 °C

Storage
Conditions

-150 -196 -80 °C

Sterility

PCR