

L-929-GFP | 305956

Description	L-929-GFP	L-929	L-929	L-929-C
	L-929-GFP	GFP		

Organism

Tissue

Synonyms	L929/GL50
-----------------	-----------

Age	100
------------	-----

Gender

Cell type

Growth properties

Citation	L929-GFP Cytion	305956
-----------------	-----------------	--------

Biosafety level	1
------------------------	---

NCBI_TaxID	10090
-------------------	-------

CellosaurusAccession	CVCL_E2Z7
-----------------------------	-----------

Culture Medium	DMEM:Ham's F12 1:1	w: 3.1 g/L	w: 2.5 mM L-	w: 15 mM HEPES	w: 0.5 mM	w: 1.2 g/L NaHCO3	Cytion	8
-----------------------	--------------------	------------	--------------	----------------	-----------	-------------------	--------	---

L-929-GFP | 305956

Supplements	10% FBS																																																																																																														
Dissociation Reagent	Accutase																																																																																																														
Subculturing	PBS	T25	3-5	PBS	T75	5-10	Accutase	T25	1-2	T75	2.5																																																																																																				
Seeding density	1 3×10^4 /																																																																																																														
Freeze medium	+10% DMSO																																																																																																														
Thawing and Culturing Cells	<table border="1"> <tr> <td>1.</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td>2.</td> <td></td> <td>-150°C</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>3</td> <td></td> <td></td> <td></td> </tr> <tr> <td>3.</td> <td></td> <td>37°C</td> <td></td> <td></td> <td></td> <td>40-60</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td>4.</td> <td></td> <td></td> <td></td> <td>70%</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td>5.</td> <td></td> <td></td> <td>8</td> <td></td> <td>15</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td>6.</td> <td>200 x g</td> <td></td> <td>5</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td>7.</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> </table>													1.														2.		-150°C								3				3.		37°C				40-60								4.				70%										5.			8		15									6.	200 x g		5											7.													
1.																																																																																																															
2.		-150°C								3																																																																																																					
3.		37°C				40-60																																																																																																									
4.				70%																																																																																																											
5.			8		15																																																																																																										
6.	200 x g		5																																																																																																												
7.																																																																																																															
Incubation Atmosphere	37°C, 5% CO ₂																																																																																																														
Shipping Conditions	-78 °C																																																																																																														
Storage Conditions	-150 -196 -80 °C																																																																																																														

/ **/HLA**