

NCI-H2126 | 300639

<b>Description</b>	NCI-H2126	NSCLC			
	NCI-H2126		6q 13q	NSCLC	
	NCI-H2126				NSCLC

**Organism**

**Tissue**

**Disease**

**Metastatic site**

**Applications**

**Synonyms** H-2126 NCI-H2126 NCI-H2126

**Age** 65

**Gender**

**Ethnicity**

**Morphology**

**Growth properties**

**Citation** NCI-H2126 Cytion 300639

**Biosafety level** 2

**NCBI\_TaxID** 9606

NCI-H2126 | 300639

**CellosaurusAccession** CVCL\_1532

**Isoenzymes** AK-1 1 ES-D 1-2 G6PD B GLO-I 2 Me-2 0 PGM1 1-2 PGM3 2

**Tumorigenic**

**Viruses** EB

**Ploidy status**

**Culture Medium** DMEM:Ham's F12 1:1 w: 3.1 g/L w: 2.5 mM L- w: 15 mM HEPES w: 0.5 mM w: 1.2 g/L NaHCO3 Cytion 8

**Supplements** 5% FBS 0.005 mg/mL 0.01 mg/mL 30nM 10 nM 10 nM beta-estradiol

**Dissociation Reagent** Accutase

**Subculturing** PBS T25 3-5 PBS T75 5-10 Accutase T25 1-2 T75 2.5

**Freeze medium** FBS +10% DMSO CM-1 Cytion 800100 CM-1 Cytion 800100

NCI-H2126 | 300639

**Thawing and Culturing Cells**

1.				
2.		-150°C		3
3.		37°C	40-60	
4.		70%		
5.		8	15	
6.	300 x g	3		
7.	10		T25	T25
8.				

**Incubation Atmosphere**

37°C, 5% CO<sub>2</sub>

**Flask Coating**

**Freezing Procedure**

-78 °C

**Shipping Conditions**

-78 °C

**Storage Conditions**

-150 -196 -80 °C

/ /HLA

**Sterility**

PCR