

NCI-H2126 | 300639

<b>Description</b>	NCI-H2126	NSCLC			
	NCI-H2126		6q 13q	NSCLC	
	NCI-H2126				NSCLC

**Organism**

**Tissue**

**Disease**

**Metastatic site**

**Applications**

**Synonyms** H-2126 NCI-H2126 NCI-H2126

**Age** 65

**Gender**

**Ethnicity**

**Morphology**

**Growth properties**

**Citation** NCI-H2126 Cytion 300639

**Biosafety level** 2

**NCBI\_TaxID** 9606

NCI-H2126 | 300639

**CellosaurusAccession** CVCL\_1532

**Isoenzymes** AK-1 1 ES-D 1-2 G6PD B GLO-I 2 Me-2 0 PGM1 1-2 PGM3 2

**Tumorigenic**

**Viruses** EB

**Ploidy status**

**Culture Medium** DMEM:Ham's F12 1:1 w: 3.1 g/L w: 2.5 mM L- w: 15 mM HEPES w: 0.5 mM w: 1.2 g/L NaHCO3 Cytion 8

**Supplements** 5% FBS 0.005 mg/mL 0.01 mg/mL 30nM 10 nM 10 nM beta-estradiol

**Dissociation Reagent** Accutase

**Subculturing** PBS T25 3-5 PBS T75 5-10 Accutase T25 1-2 T75 2.5

**Freeze medium** FBS +10% DMSO CM-1 Cytion 800100 CM-1 Cytion 800100

NCI-H2126 | 300639

**Thawing and  
Culturing Cells**

- 1.
2. -150°C 3
3. 37°C 40-60
4. 70%
5. 8 15
6. 300 x g 3
7. 10 T25 T25
- 8.

**Incubation  
Atmosphere**

37°C, 5% CO<sub>2</sub>

**Flask Coating**

**Freezing  
Procedure**

-78 °C

**Shipping  
Conditions**

-78 °C

**Storage  
Conditions**

-150 -196 -80 °C

/ /HLA

**Sterility**

PCR