

C127I | 400134

<b>Description</b>	C127I	BALB/c	C127I
	C127I		C127I
<b>Organism</b>			
<b>Tissue</b>			
<b>Disease</b>			
<b>Applications</b>	DNA		
<b>Synonyms</b>	C 127I C-127I C-127 I CNC 127I		
<b>Breed/Subspecies</b>	RIII		
<b>Gender</b>			
<b>Morphology</b>			
<b>Growth properties</b>			
<b>Citation</b>	C127I Cytion 400134		
<b>Biosafety level</b>	1		
<b>NCBI_TaxID</b>	10090		
<b>CellosaurusAccession</b>	CVCL_3882		
<b>GMO Status</b>	GMO-S1 C127I T7 RNA CFTR		C127

C127I | 400134

**Viruses**

**Virus susceptibility**

**Reverse transcriptase**

**Culture Medium** DMEM w 4.5 / w 4 L- w 3.7 / NaHCO3 w 1.0 Cytion 820300a

**Supplements** 10% FBS

**Dissociation Reagent** Accutase

**Subculturing** PBS T25 3-5 PBS T75 5-10 Accutase T25 1-2 T75 2.5

**Fluid renewal** 2 3

**Post-Thaw Recovery** <sup>4</sup> /5×10 24

**Freeze medium** FBS +10% DMSO CM-1 Cytion 800100 CM-1 Cytion 800100

C127I | 400134

**Thawing and Culturing Cells**

- 1.
2. -150°C 3
3. 37°C 40-60
4. 70%
5. 8 15
6. 300 x g 3
7. 10 T25 T25
- 8.

**Incubation Atmosphere** 37°C, 5% CO<sub>2</sub>

**Flask Coating**

**Freezing Procedure** -78 °C

**Shipping Conditions** -78 °C

**Storage Conditions** -150 -196 -80 °C

/ /HLA

**Sterility** PCR