

AH-130 | 500412

**Description** Yoshida Yoshida 1956 AH-130

**Organism**

**Tissue**

**Disease**

**Metastatic site**

**Synonyms** AH-130 AH130 AH130 AH 130 AH-130 AH130-TC AH130/P

**Breed/Subspecies** -

**Morphology**

**Growth properties** /

**Citation** AH-130 Cytion 500412

**Biosafety level** 1

**NCBI\_TaxID** 10116

**CellosaurusAccession** CVCL\_4367

**Tumorigenic** Wistar

**Viruses** RAP .

AH-130 | 500412

**Virus susceptibility**

**Culture Medium** DMEM:Ham's F12 1:1 w: 3.1 g/L w: 2.5 mM L- w: 15 mM HEPES w: 0.5 mM w: 1.2 g/L NaHCO3 Cytion 8

**Supplements** 10% FBS

**Dissociation Reagent** Accutase

**Subculturing** 15 PBS T25 3-5 T75 5-10 Accutase T25 1-2 T75 2.5

**Seeding density**  $2 \times 10^4 /$

**Fluid renewal** 3 5

**Post-Thaw Recovery**

**Freeze medium** FBS +10% DMSO CM-1 Cytion 800100 CM-1 Cytion 800100

AH-130 | 500412

Thawing and Culturing Cells				
1.				
2.		-150°C		3
3.		37°C	40-60	
4.		70%		
5.		8	15	
6.	300 x g	3		
7.	10		T25	T25
8.				

**Incubation Atmosphere** 37°C, 5% CO<sub>2</sub>

**Flask Coating**

**Freezing Procedure** -78 °C

**Shipping Conditions** -78 °C

**Storage Conditions** -150 -196 -80 °C

/ /HLA

**Sterility** PCR