

HPAF-II | 305088

Description	HPAF-II				HPAF-II
	HPAF-II		KRAS	TP53	CDKN2A

Organism

Tissue

Disease

Metastatic site

Synonyms HPAF II HPAFII HPAF-2 HPAF2 HPAF/CD18 CD18/HPAF HPAF-II/CD18 CD-18 CD18 CD 18

Age 44

Gender

Ethnicity

Morphology

Growth properties

Citation HPAF-II Cytion 305088

Biosafety level 1

NCBI_TaxID 9606

CellosaurusAccession CVCL_0313

HPAF-II | 305088

Culture Medium EMEM MEM Eagle w 2 mM L- w 2.2 g/L NaHCO3 w EBSS Cytion 820100a

Supplements 10% FBS 1% NEAA

Dissociation Reagent Accutase

Doubling time 26

Subculturing PBS T25 3-5 PBS T75 5-10 Accutase T25 1-2 T75 2.5

Fluid renewal 2 3

Freeze medium FBS +10% DMSO CM-1 Cytion 800100 CM-1 Cytion 800100

Thawing and Culturing Cells

- 1.
2. -150°C 3
3. 37°C 40-60
4. 70%
5. 8 15
6. 300 x g 3
7. 10 T25 T25
- 8.

Incubation Atmosphere 37°C, 5% CO₂

HPAF-II | 305088

Flask Coating

**Freezing
Procedure**

-78 °C

**Shipping
Conditions**

-78 °C

**Storage
Conditions**

-150 -196 -80 °C

/ /HLA

Sterility

PCR