

SK-N-BE(2) | 305058

Description β ⁶- / SK-N-BE(2) 1×10

Organism

Tissue

Disease

Metastatic site

Synonyms SK-N-BE2 SK-N-BE-2 SKNBE 2 SKNBE-2 SKNBE2 SK-N-BE SKNBE

Age 2

Gender

Ethnicity

Morphology

Growth properties /

Citation SK-N-BE(2) Cytion 305058

Biosafety level 1

NCBI_TaxID 9606

CellosaurusAccession CVCL_0528

Tumorigenic

SK-N-BE(2) | 305058

Culture Medium	50:50	EMEM	Ham's F12	Cytion	820100a	820600a
-----------------------	-------	------	-----------	--------	---------	---------

Supplements	10% FBS
--------------------	---------

Dissociation Reagent	Accutase
-----------------------------	----------

Subculturing	15	PBS	T25	3-5	T75	5-10	Accutase	T25	1-2	T75	2.5
---------------------	----	-----	-----	-----	-----	------	----------	-----	-----	-----	-----

Fluid renewal	2	3
----------------------	---	---

Freeze medium	50%	+ 40% FBS + 10% DMSO	CM-1	Cytion	800100
----------------------	-----	----------------------	------	--------	--------

Thawing and Culturing Cells	1.					
	2.		-150°C			3
	3.		37°C		40-60	
	4.			70%		
	5.		8	15		
	6.	300 x g	3			
	7.	10			T25	T25
	8.					

Incubation Atmosphere	37°C, 5% CO ₂
------------------------------	--------------------------

Flask Coating

SK-N-BE(2) | 305058

Freezing Procedure -78 °C

Shipping Conditions -78 °C

Storage Conditions -150 -196 -80 °C

/ **/HLA**

Sterility PCR