

LXF-289 | 300269

| | | | | | |
|--------------------|---------|----|--------------|----------|---------|
| Description | LxF-289 | 63 | 50 | LxF-289 | NSCLC |
| | LxF-289 | | | 70 Hsp70 | LxF-289 |
| | LxF-289 | | S- GSTπ mRNA | | |

Organism

Tissue

Disease

Synonyms LxF289 LxF 289 LxF 289L

Age 62

Gender

Ethnicity

Morphology

Growth properties

Citation LxF-289 Cytion 300269

Biosafety level 1

NCBI_TaxID 9606

CellosaurusAccession CVCL_1394

LXF-289 | 300269

Tumorigenic

Reverse transcriptase

Culture Medium RPMI 1640 w 2.0 mM w 2.0 g/L NaHCO₃ Cytion 820700a

Supplements 10% FBS

Dissociation Reagent Accutase

Subculturing PBS T25 3-5 PBS T75 5-10 Accutase T25 1-2 T75 2.5

Seeding density 1×10^4 /

Fluid renewal 3 5

Post-Thaw Recovery 24 48

Freeze medium FBS +10% DMSO CM-1 Cytion 800100 CM-1 Cytion 800100

LXF-289 | 300269

| Thawing and Culturing Cells | | | | |
|-----------------------------|---------|--------|-------|-----|
| 1. | | | | |
| 2. | | -150°C | | 3 |
| 3. | | 37°C | 40-60 | |
| 4. | | | 70% | |
| 5. | | 8 | 15 | |
| 6. | 300 x g | 3 | | |
| 7. | 10 | | T25 | T25 |
| 8. | | | | |

Incubation Atmosphere 37°C, 5% CO₂

Flask Coating

Freezing Procedure -78 °C

Shipping Conditions -78 °C

Storage Conditions -150 -196 -80 °C

/ /HLA

Sterility PCR